



Media Name	Media Type	Organism Cultured	Common Brewery Organism	Notes
Universal Beer Agar	Aerobic or Anaerobic	Brewer's yeast, wild yeast, bacteria, molds	<i>Brettanomyces, Candida, Saccharomyces</i> -type wild yeast, <i>Lactobacillus, Bacillus, Acetobacter</i>	Cycloheximide can be added to inhibit brewer's yeast
Wallerstein Differential (WLD)	Aerobic or Anaerobic	Wild yeast , bacteria. mold	<i>Brettanomyces, Candida, Saccharomyces</i> -type wild yeast, <i>Lactobacillus, Bacillus, Acetobacter</i>	
SDA/LMDA	Aerobic or Anaerobic	Bacteria	Acetic acid bacteria, <i>Bacillus, Lactobacillus, Enterobacter, Pediococcus</i>	Cycloheximide can be added to inhibit brewer's yeast
Hsu's Lactobacillus and Pediococcus (HLP)	Anaerobic	Bacteria (Occasionally wild yeast)	<i>Lactobacillus and Pediococcus</i>	
Lin's Wild Yeast Medium (LWYM)	Aerobic	Wild Yeast	<i>Saccharomyces</i> -type wild yeast	Some brewer's strains will grow
Lin's Cupric Sulfate Medium (LCSM)	Aerobic	Wild Yeast	Non- <i>Saccharomyces</i> wild yeast	Some brewer's strains will grow
Wallerstein Nutrient Media (WLN)	Aerobic	Brewer's yeast, wild yeast, bacteria, molds	Brewer's yeast , wild yeast, <i>Lactobacillus, Bacillus, Acetobacter</i>	